Circulant Distant Two Labeling and Circular Chromatic Number

Daphne Der-Fen Liu *
Department of Mathematics and Computer Science
California State University, Los Angeles
Los Angeles, CA 90032, USA
Emil: dliu@calstatela.edu

Xuding Zhu †
Department of Applied Mathematics
National Sun Yat-sen University
Kaoshing, Taiwan 80424
Email: zhu@ibm18.math.nsysu.edu.tw

May 25, 2000

Abstract

Let G be a graph and d, d' be positive integers, $d' \geq d$. An $m-(d,d')-$circular distance two labeling is a function $f : V(G) \to \{0, 1, 2, \cdots, m-1\}$ such that $|f(u) - f(v)|_m \geq d$ if u and v are adjacent; and $|f(u) - f(v)|_m \geq d'$ if u and v are distance two apart, where $|x|_m := \min\{|x|, m - |x|\}$. The minimum m such that there exists an $m-(d,d')-$circular labeling for G is called the $\sigma_{d,d'}$-number of G and denoted by $\sigma_{d,d'}(G)$. The $\sigma_{d,d'}$-numbers for trees can be obtained by a first-fit algorithm. In this article, we completely determine the $\sigma_{d,1}$-numbers for cycles. In addition, we show connections between generalized circular distance labeling and circular chromatic number.

Keywords. Vertex-labeling, distance two labeling, circular chromatic number.

*Supported in part by the National Science Foundation under grant DMS 9805945.
†Supported in part by the National Science Council, R. O. C., under grant 88-2115-M-110-001.

1 Introduction

For a graph G, the distance between any two vertices u and v is denoted by $\mathrm{d}_G(u,v)$. Given G and positive integers d, d' with $d \geq d'$, an $L(d,d')-$labeling (distance two labeling) of G is a function $f : V(G) \to \{0,1,2,\cdots\}$ such that $|f(u) - f(v)| \geq d$ if $uv \in E(G)$; and $|f(u) - f(v)| \geq d'$ if $\mathrm{d}_G(u,v) = 2$. The *span* of f is defined as $\min_{u,v\in V(G)} |f(u)-f(v)|$. The minimum span of an $L(d,d')-$labeling for G is denoted by $\lambda_{d,d'}(G)$. The $L(d,d')-$labelings, for different values of d and d', have been studied by several authors in the past decade. (See [1, 2, 3, 4, 8, 10, 12].)

The circular distance two labeling is a variation of $L(d,d')-$labeling by using a different measurement. Given G and positive integers d and d' with $d \geq d'$, an $m-(d,d')_c-$ labeling is a function, $f : V(G) \to \{0,1,2,\cdots,m-1\}$, such that $|f(u) - f(v)|_m \geq d$ if $uv \in E(G)$, and $|f(u) - f(v)|_m \geq d'$ if $\mathrm{d}_G(u,v) = 2$, where $|x|_m := \min\{|x|, m-|x|\}$. For any graph G and any $d \geq d'$, the $\sigma_{d,d'}-$*number* of G, $\sigma_{d,d'}(G)$, is the minimum m such that there exists an $m-(d,d')_c-$labeling for G.

In Section 2, we give exact values of $\sigma_{d,d'}-$number for trees and $\sigma_{d,1}-$number for cycles. Georges and Mauro [3] proved that $\lambda_{ad,ad'}(G) = a\lambda_{d,d'}(G)$ holds for any graph G and any positive integers a, d, d' with $d \geq d'$. The results for cycles and trees shown in this article indicate that a similar result of Georges and Mauro does not hold for circular distance two labelings. That is, there exist graphs such that $\sigma_{ad,ad'}(G) \neq a\sigma_{d,d'}(G)$ for some positive integer a.

In Section 3, we investigate close relations between generalized circular distance labeling and circular chromatic number.

2 The σ-number for trees and cycles

In this section, we give the exact values of the $\sigma_{d,d'}$-numbers for trees for all $d \geq d'$, and $\sigma_{d,1}$-numbers for cycles.

Theorem 2.1 *Let T be a tree with maximum degree Δ. Then for any $d \geq d'$, $\sigma_{d,d'}(T) = 2d + (\Delta - 1)d'$.*

Proof. Observe that for any star $G = K_{1,n}$, $\sigma_{d,d'}(G) = 2d + (n-1)d'$. Therefore, $\sigma_{d,d'}(T) \geq 2d + (\Delta - 1)d'$, since T contains a $K_{1,\Delta}$. So it suffices to find a $(2d + (\Delta - 1)d')$-$(d, d')_c-$labeling for T. This can be done by a first-fit algorithm. First, fix a root of T of degree Δ, label it by 0 and its neighbors by $d, d+d', d+2d', \cdots, d+(\Delta-1)d'$; then label the neighbors of these labeled vertices one by one using the labels from the set $\{0, 1, 2, \cdots, 2d + (\Delta-1)d' - 1\}$; and continue this process until all vertices are labeled. At each step, if a vertex v is labeled by x, then there are exactly Δ labels, $\{x + d, x + d + d', x + d + 2d', \cdots, x + d + (\Delta - 1)d'\} \pmod{2d + (\Delta - 1)d'}$, that can be used by the neighbors of v. Thus, this process produces a valid $(2d + (\Delta - 1)d')$-$(d, d')_c-$labeling for T. Q.E.D.

Theorem 2.2 *For any $d \geq d'$, $\sigma_{d,d'}(C_n) \geq 2d + d'$. Furthermore, $\sigma_{d,d'}(C_n) = 2d + d'$ if and only if $n \equiv 0 \pmod{\frac{2d+d'}{r}}$, where $r = gcd\ (2d + d', d)$.*

Proof. Observe that $\sigma_{d,d'}(C_n) \geq 2d + d'$, since $\sigma_{d,d'}(K_{1,2}) = 2d + d'$. Suppose $\sigma_{d,d'}(C_n) = 2d + d'$ and let f be a $(2d + d') - (d, d')_c-$labeling for C_n. For any vertex v of C_n, suppose $f(v) = x$ for some $0 \leq x \leq 2d + d' - 1$, then there are exact two values, $x - d$, $x + d \pmod{2d + d' - 1}$, that can be assigned to the two neighbors of x. Let the vertices of C_n, in clockwise order, be $v_1, v_2, \cdots, v_n$. Without loss of generality, assume $f(v_1) = 0$ and $f(v_2) = d$. Then $f(v_{i+1}) = (f(v_i) + d) \bmod (2d + d')$. Since $f(v_n)$ must be $d + d'$, f is well-defined only when $n \equiv 0 \pmod{\frac{2d+d'}{r}}$, where $r = \gcd(2d + d', d)$. Q.E.D.

Corollary 2.3 *Suppose $d \geq d'$, then $\sigma_{kd,kd'}(C_n) = k\sigma_{d,d'}(C_n) = k(2d + d')$ if $n \equiv 0 \pmod{\frac{2d+d'}{r}}$, where $r = gcd(2d + d', d)$.*

It is known [7] and not hard to observe the following inequalities:

$$\lambda_{d,d'}(G)+1 \le \sigma_{d,d'}(G) \le \lambda_{d,d'}(G)+d, \quad \text{for any graph } G. \qquad (*)$$

To derive the values of $\sigma_{d,1}(C_n)$, we first quote the result of Georges and Mauro [3] on $\lambda_{d,1}(C_n)$.

Theorem 2.4 *([3]) Let C_n be a cycle, then*

$$\lambda_{d,1}(C_n) = \begin{cases} d+2, & n \equiv 0 \pmod 4; \\ d+3, & n \equiv 2 \pmod 4; \\ 2d, & n \equiv 1 \pmod 2. \end{cases}$$

Combining (*) with Theorems 2.2 and 2.4. we have

$$2d+1 \le \sigma_{d,1}(C_n) \le \begin{cases} 2d+2, & n \equiv 0 \pmod 4; \\ 2d+3, & n \equiv 2 \pmod 4; \\ 3d, & n \equiv 1 \pmod 2. \end{cases}$$

Theorem 2.5 *Let C_n be a cycle and let $d \ge 2$. Then $\sigma_{d,1}(C_3) = 3d$; and for $n \ge 4$,*

$$\sigma_{d,1}(C_n) = \begin{cases} 2d+1, & \text{if } n \equiv 0 \pmod{2d+1}; \\ 2d+2, & \text{if } n \text{ is even, and } n \not\equiv 0 \pmod{2d+1}; \\ 2d+2, & \text{if } n \text{ is odd, } d < n, \text{ and } n \not\equiv 0 \pmod{2d+1}; \\ 2d+\lceil \frac{d}{k} \rceil, & \text{if } n \text{ is odd, } n = 2k+1,\ d \ge n, \text{ and } n \not\equiv 0 \pmod{2d+1}. \end{cases}$$

Proof. It is easy to see that $\sigma_{d,1}(C_3) = 3d$. For $n \ge 4$, because gcd $(2d+1, d) = 1$, by Theorem 2.2, $\sigma_{d,1}(C_n) \ge 2d+1$, and $\sigma_{d,1}(C_n) = 2d+1$ if and only if $n \equiv 0 \pmod{2d+1}$. This proves the first case in the formula.

To complete the proof, it suffices to claim the following cases:

Case 1 n is even and $n \not\equiv 0 \pmod{2d+1}$. It suffices to find a $(2d+2)-(d,1)_c$−labeling. Let $n = 2k$ and let the vertices of C_{2k}, in clockwise order, be v_1, v_2, v_3, $\cdots$, v_{2k}. Define the labeling f by $f(v_1) = 0$; and

$$f(v_{i+1}) = \begin{cases} (f(v_i)+d) \bmod (2d+2), & \text{if } 1 \le i \le k-1; \\ (f(v_i)+d+2) \bmod (2d+2), & \text{if } k+1 \le i \le 2k-1; \end{cases}$$

and $f(v_{k+1}) = (f(v_k)+d+1) \bmod (2d+2)$. Then $f(v_{2k}) = d+1$ (since $f(v_{2k}) = (2k-1)d+2k-1 = (2d+2)k-(d+1) \equiv d+1 \pmod{2d+2}$). Moreover, f is a

$(2d+2)-(d,1)_c$−labeling for C_n, since the circular difference (mod $2d+2$) between any adjacent vertices is either d or $d+1$, and any pair of vertices of distance two receive different labels.

Case 2 n is odd, $n = 2k+1$, $n \geq 2$, $d < n$, and $n \not\equiv 0 \pmod{2d+1}$. Again it suffices to find a $(2d+2)-(d,1)_c$−labeling. Let the vertices of C_{2k+1}, in clockwise order, be $v_1, v_2, \cdots, v_{2k+1}$. By considering the parity of d, we separate this case into two subcases:

Case 2.1 d is even. Let $d = 2m$, and $x = k - m \geq 0$. If $x = 0$, define the coloring f by $f(v_1) = 0$; and $f(v_{i+1}) = (f(v_i) + d) \bmod (2d+2)$ for all $1 \leq i \leq 2k$. Then $f(v_{2k+1}) = d+2$ (since $f(v_{2k+1}) = 2kd = k(2d+2) - 2k = k(2d+2) - d \equiv d+2 \pmod{2d+2}$). Hence f is a $(2d+2)-(d,1)_c$−labeling for C_n.

If $x = 1$, define the coloring f by $f(v_1) = 0$; and

$$f(v_{i+1}) = \begin{cases} (f(v_i)+d+1) \bmod (2d+2), & \text{if } i = 1, 3; \\ (f(v_i)+d+2) \bmod (2d+2), & \text{if } i = 2 \text{ and } 4 \leq i \leq 2k. \end{cases}$$

Then $f(v_{2k+1}) = d+2$, and f is a $(2d+2)-(d,1)_c$−labeling for C_n.

If $x > 1$ define the coloring f by $f(v_1) = 0$; and

$$f(v_{i+1}) = \begin{cases} (f(v_i)+d+1) \bmod (2d+2), & \text{if } v = 1, x+1; \\ (f(v_i)+d+2) \bmod (2d+2), & \text{if } 2 \leq i \leq x; \\ (f(v_i)+d) \bmod (2d+2), & \text{if } x+2 \leq i \leq 2k. \end{cases}$$

Then $f(v_{2k+1}) = d+2 \pmod{2d+1}$ and f is a $(2d+2)-(d,1)_c$−labeling for C_n.

Case 2.2 d is odd. Let $d = 2m+1$. Then $k > m$. Let $x = k - m \geq 1$. If $x = 1$, define the coloring f by $f(v_1) = 0$; $f(v_2) = d+1$; and $f(v_{i+1}) = (f(v_i) + d) \bmod (2d+2)$ for $2 \leq i \leq 2k$. Then $f(v_{2k+1}) = d+2$, and f is a $(2d+2)-(d,1)_c$−labeling for C_n.

If $x = 2$, define the coloring f by $f(v_1) = 0$; and

$$f(v_{i+1}) = \begin{cases} (f(v_i)+d+1) \bmod (2d+2), & \text{if } i = 1, 3, 5; \\ (f(v_i)+d) \bmod (2d+2), & \text{if } i = 2, 4 \text{ or } 6 \leq i \leq 2k. \end{cases}$$

Because $k = m+2 \geq 3$, so f is well-defined. Furthermore, it is easy to verify that $f(v_{2k+1}) = d+2$, and f is a $(2d+2)-(d,1)_c$−labeling for C_n.

If $x > 2$, define the coloring f by $f(v_1) = 0$; and

$$f(v_{i+1}) = \begin{cases} (f(v_i) + d + 1) \bmod (2d+2), & \text{if } i = 1, x, 2k; \\ (f(v_i) + d + 2) \bmod (2d+2), & \text{if } 2 \le i \le x-1; \\ (f(v_i) + d) \bmod (2d+2), & \text{if } x+1 \le i \le 2k-1. \end{cases}$$

Then $f(v_{2k+1}) = d+2$, and f is a $(2d+2) - (d,1)_c$–labeling for C_n.

Case 3 n is odd, $n = 2k+1$, $d \ge n$, and $n \not\equiv 0 \pmod{2d+1}$. Let $\lceil \frac{d}{k} \rceil = q$. Suppose $\sigma_{d,1}(C_{2k+1}) = l \le 2d + q - 1$. Let f be an $l - (d,1)_c$–labeling for C_{2k+1}. Let the vertices of C_{2k+1}, in clockwise order, be v_0, a_1, a_2, $\cdots$, a_k, b_k, $b_{k-1}, \cdots, b_1$. Without loss of generality, let $f(v_0) = 0$. By definition of circular labeling, we have

$$\begin{array}{ll} & |f(a_1) - f(b_1)|_l \le l - 2d \le q - 1 \\ \Rightarrow & |f(a_2) - f(b_2)|_l \le 2(q-1) \\ \Rightarrow & |f(a_3) - f(b_3)|_l \le 3(q-1) \\ \Rightarrow & \cdots\cdots\cdots \\ \Rightarrow & |f(a_k) - f(b_k)|_l \le k(q-1) < d. \end{array}$$

The last inequality is true since $\lceil \frac{d}{k} \rceil = q$. This contradicts the fact that a_k and b_k are adjacent.

To complete the proof, it is enough to find a $(2d+q) - (d,1)_c$–labeling for C_{2k+1}. Now we let the vertices of C_n, in clockwise order, be $v_1, v_2, \cdots, v_{2k+1}$ and let $d = xk + r$ for some x and $1 \le r \le k$. Then $q = x+1$. If $r = k$, define the coloring f by $f(v_1) = 0$; and $f(v_{i+1}) = (f(v_i) + d) \bmod (2d+q)$, $1 \le i \le 2k$. Then $f(v_{2k+1}) = d + q$ (since $f(v_{2k+1}) = 2kd = k(2d+q) - kq \equiv -d \equiv d + q \pmod{2d+q}$). Hence f is a $(2d+q) - (d,1)_c$–labeling for C_n.

If $1 \le r < k$, define the coloring by $f(v_1) = 0$; and

$$f(v_{i+1}) = \begin{cases} (f(v_i) + d + 1) \bmod (2d+q), & \text{if } i = 1, 3, 5, \cdots, 2(k-r) - 1; \\ (f(v_i) + d) \bmod (2d+q), & \text{otherwise.} \end{cases}$$

Then $f(v_{2k+1}) = d + q$, and f is a $(2d+q) - (d,1)_c$–labeling for C_n. Q.E.D.

The result above has also been proved, lately and independently, by Wu and Yeh [13].

3 Generalized Circular Distance Labeling and Its Relations to Circular Chromatic Number

The circular distance two labeling is a special case of circular distance p labeling, $p \geq 1$, introduced and studied by ven den Heuvel at el [7]. Given a graph G and integers $d_1 \geq d_2 \geq \cdots \geq d_p \geq 1$, an $m-$labeling f of G, $f : V(G) \rightarrow \{0, 1, 2, \cdots, m-1\}$, satisfies the constraints $(d_1, d_2, \cdots, d_p)$ if $|f(u) - f(v)|_m \geq d_i$, for all $i \in \{1, 2, \cdots, p\}$ and $\mathrm{d}_G(u, v) = i$. The minimum m of such a labeling for G is denoted by $\sigma(G; d_1, d_2, \cdots, d_p)$; and for the case $d_1 = d_2 = \cdots d_p$, it is denoted by $\sigma(G; (d_1)^p)$.

The σ-number of a graph G is closely related to the circular chromatic number of G. Let a and b be positive integers such that $a \geq 2b$. An (a, b)-coloring of a graph $G = (V, E)$ is a mapping c from V to $\{0, 1, \cdots, a-1\}$ such that $|c(x) - c(y)|_a \geq b$ for any edge xy in G. The *circular chromatic number* $\chi_c(G)$ of G is the infimum of a/b for which there exists an (a, b)-coloring of G. The circular chromatic number is also known as the *star-chromatic number* in the literature [11].

Theorem 3.1 *For any graph G and positive integer d, $\sigma(G; d) = \lceil d\chi_c(G) \rceil$.*

Proof. By definition, $\chi_c(G) \leq \frac{\sigma(G;d)}{d}$. Since $\sigma(G; d)$ is an integer, we have $\sigma(G; d) \geq \lceil d\chi_c(G) \rceil$.

Let $\chi_c(G) = p/q$, $\gcd(p, q) = 1$, and $m = \lceil pd/q \rceil$. To complete the proof, it suffices to find an $m-$labeling f, $f : V(G) \rightarrow \{0, 1, 2, \cdots, m-1\}$ such that $|f(u) - f(v)|_m \geq d$ for all $uv \in E(G)$. This is equivalent to finding an $(m, d)-$coloring for G. Since $\chi_c(G) = p/q$, there is a (p, q)-coloring for G, implying the existence of an $(m, d)-$coloring for G. Q.E.D.

Given a graph G and a positive integer k, the *k-th power of G*, denoted by G^k, is defined by $V(G^k) = V(G)$ and $uv \in E(G^k)$ if and only if $\mathrm{d}_G(u, v) \leq k$. It is easy to see that $\sigma(G; (d)^k) = \sigma(G^k; d)$, so we have:

Corollary 3.2 *For any graph G and positive integers k and d, $\sigma(G;(d)^k) = \lceil d\chi_c(G^k) \rceil$.*

It is known [9] that $\chi_c(C_n^k) = \frac{n}{\lfloor \frac{n}{k+1} \rfloor}$, thus we have:

Corollary 3.3 *For any positive integers $n \geq 3$, k and d, $\sigma(C_n;(d)^k) = \lceil \frac{nd}{\lfloor \frac{n}{k+1} \rfloor} \rceil$.*

References

[1] G. J. Chang and D. Kuo, *The $L(2,1)-$labeling problem on graphs*, SIAM J. Disc. Math. 9 (1996) 309–316.

[2] G. J. Chang, W. Ke, D. Kuo, D. D.-F. Liu and R. Yeh, *On $L(d,1)-$labelings of graphs*, Disc. Math., to appear.

[3] J. P. Georges and D. W. Mauro, *Generalized vertex labelings with a condition at distance two*, Congr. Numer. 109 (1995) 141–159.

[4] J. Georges, D. Mauro and M. Whittlesey, *Relating path covering to vertex labelings with a condition at distance two*, Disc. Math. 135 (1994) 103–111.

[5] J. R. Griggs and R. K. Yeh, *Labeling graphs with a condition at distance 2*, SIAM J. Discrete Math. 5 (1992) 586–595.

[6] W. K. Hale, *Frequency assignment: Theory and applications*, Proc. IEEE, 68 (1980) 1497–1514.

[7] J. van den Heuvel, R. A. Leese and M. A. Shepherd, *Graph labelling and radio channel assignment*, J. Graph Theory 29 (1998) 263–283.

[8] K. Jonas, *Graph coloring analogous with a condition at distance two: $L(2,1)-$labelings and list $\lambda-$labelings*, Ph.D. Thesis, Dept. of Math., Univ. of South Carolina, Columbia, SC, 1993.

[9] G. Gao and X. Zhu, *Star-extremal graphs and the lexicographic product*, Disc. Math. 152 (1996) 147-156.

[10] D. D.-F. Liu and R. K. Yeh, *On distance two labellings of graphs*, Ars Comb. 47 (1997) 13-22.

[11] A. Vince, *Star chromatic number*, J. Graph Theory, 12 (1988), pp. 551-559.

[12] M. A. Whittlesey, J. P. Georges and D. W. Mauro, *On the λ-number of Q_n and related graphs*, SIAM J. Disc. Math. 8 (1995) 499–506.

[13] K.-F. Wu and R. Yeh *Labelling Graphs with the Circular Difference*, manuscript.